

A GIANT GLITCH IN THE ENERGETIC 69 MS X-RAY PULSAR AXS J161730–505505

KEN'ICHI TORII¹, E.V. GOTTHELF², G. VASISHT³, T. DOTANI⁴, & K. KINUGASA⁵

Draft of Mar 10 2000

ABSTRACT

We present new results on the recently discovered 69 ms X-ray pulsar AXS J161730–505505, the sixth youngest sample of all known pulsars. We have undertaken a comprehensive X-ray observing campaign of AXS J161730–505505 with the *ASCA*, *SAX*, and *XTE* observatories and follow its long term spin-down history between 1989 and 1999, using these, archival *GINGA* and *ASCA* data sets, and the radio ephemeris. The spin-down is not simply described by a linear function as originally thought, but instead we find evidence of a giant glitch ($|\Delta P/P| \gtrsim 10^{-6}$) between 1993 August and 1997 September, perhaps the largest yet observed from a young pulsar. The glitch is well described by steps in P and $\dot{P}$ accompanied by a persistent $\ddot{P}$ similar to those in the Vela pulsar. The pulse profile of AXS J161730–505505 presents a single asymmetric peak which is maintained over all observation epochs. The energy spectrum is also steady over time, characterized by a highly absorbed power-law with a photon index $\Gamma = 1.4 \pm 0.2$, consistent with that found for other young rotation powered pulsars.

Subject headings: pulsars: general — pulsars: individual (AXS J161730–505505, PSR J1617–5055) — X-rays: general — supernova remnant — stars: neutron

1. INTRODUCTION

Radio pulsars are thought to be highly magnetized ($\sim 10^{12}$ G), rapidly spinning neutron stars whose luminosity is powered by rotational energy loss. The study of young ($< 10^5$ yrs) rotation-powered pulsars provides an important laboratory for understanding the early evolution (thermal, spin, and magnetic) of these embers of stellar collapse. The most energetic of these pulsars are observable at X-ray wavelengths, which allow us to probe these extreme, but rare, examples.

In addition to uniform spin-down corresponding to the rotational energy loss, rotation powered pulsars show sudden discontinuities in their rotation periods (see Lyne & Graham-Smith 1998, chapter 6). These rare phenomena, known as “glitches” are considered to arise from sudden changes in the configuration of super-dense material in the neutron star interior. To date, a total of 71 glitches with $|\Delta P/P| > 10^{-9}$ have been reported in 30 pulsars (Urama & Okeke 1999). Observation of pulsar glitches gives us insights into the structure and physical processes inside the neutron stars, such as the interactions of neutron superfluid and crust components (e.g., Anderson & Itoh 1975).

In this article we report the detection of a giant glitch from AXS J161730–505505 by using the newly acquired multi-mission X-ray data. This source is an unusual case of a young pulsar discovered first by its X-ray emission, revealed during the course of an archival X-ray study of the SNR RCW 103 (Gotthel, Petre, & Hwang 1997). Further analysis detected highly significant pulsations from photons attributed to this source (Torii *et al.* 1998). The 69 ms pulse period was consistent with that reported from a *GINGA* observation of the region (Aoki, Dotani, & Mitsuda 1992). Recent radio observation has confirmed AXS J161730–505505 as a young energetic rotation-powered pulsar (Kaspi *et al.* 1998). The lack of evidence for

this object in archival soft X-ray images of this field suggested that the source might be highly absorbed, making it difficult to determine whether it is associated with a SNR.

2. OBSERVATIONS

A set of day-long X-ray observations of the field containing RCW 103, 1E 161348–5055, and AXS J161730–505505 were performed with the *ASCA* (Tanaka, Inoue, & Holt 1994), *SAX* (Boella *et al.* 1997), and *XTE* (Bradt *et al.* 1993) observatories. We summarize in table 1 the set of observations presented in this work.

Imaging data were acquired with the Gas Imaging Spectrometers (GIS) on-board *ASCA* and with the Medium Energy Concentrators (MECS) instruments on *SAX*. These instruments have moderate imaging ($\sim 2'$) and spectral resolution ($\sim 8\%$ at 6 keV) over an energy band pass of 0.7–10 keV (GIS) and 1.5–12 keV (MECS), with a field-of-view (FOV) large enough to cover the SNR and pulsar simultaneously. Non-imaging data were obtained with the Proportional Counter Array (PCA) on-board *XTE* which provides broader energy band-pass (2–60 keV) at lower spectral resolution ($\sim 16\%$ at 6 keV). The PCA FOV is roughly circular with a $\sim 1^\circ$ FWHM response. The GIS data were collected in the highest time resolution mode (0.5 ms or 61 μ s, depending on data acquisition mode) whose measured absolute accuracy is 200 μ s in this mode (Saito *et al.* 1997). The PCA data were collected using the Good Xenon mode with 0.9 μ s timing resolution. For the current analysis, the absolute timing uncertainty is $\sim 100 \mu$ s (Rots *et al.* 1998). Photons collected by the MECS are time tagged with 15 μ s resolution. We do not include data from *ASCA*'s Solid-state Imaging Spectrometers (SIS), as the pulsar fell just off the edge of its FOV. Nor do we include data from *SAX*'s other instruments as

¹Space Utilization Research Program (SURP), Tsukuba Space Center (TKSC), National Space Develop Agency of Japan (NASDA), 2-1-1 Sengen, Tsukuba, Ibaraki 305-8505, Japan; torii.kenichi@nasda.go.jp

²Columbia Astrophysics Laboratory, 550 West 120th Street, New York, NY 10027; evg@astro.columbia.edu

³Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA, 91109; gv@astro.caltech.edu

⁴Institute of Space and Astronautical Science, 3-1-1 Yoshinodai, Sagami-hara, Kanagawa 229-8510, Japan; dotani@astro.isas.ac.jp

⁵Gunma Astronomical Observatory, 6860-86 Nakayama, Takayama, Agatsuma, Gunma 377-0702, Japan; kinugasa@astron.pref.gunma.jp

the observing time is insufficient for these instruments to measure the pulsar periodicity reliably, as a part of the observation was interrupted prematurely.

Each data set was processed through its standard pipeline reduction for that mission and edited to exclude times of high background contamination using the standard screening criteria. This rejects time intervals of South Atlantic Anomaly passages, Earth occultations, bright Earth limb in the FOV (ASCA and SAX only), and other periods of high particle activity. The resulting effective observation times are summarized in Table 1. For each observation, event data from all detectors were co-added and the arrival times of each event were corrected to the solar system barycenter using the software TIMECONV (ASCA), BARYCONV (SAX), or FXBARY (XTE).

3. RESULTS

3.1. Timing

The X-ray images obtained with both ASCA and SAX above 3 keV reveal AXS J161730–505505 4' outside the SNR shell (see figure 1a and 1b of Gotthelf *et al.* 1997; Gotthelf, Petre, & Vasisht 1999). To increase the signal-to-noise ratio for detecting pulsations from the pulsar, we extracted photons from an 8' diameter aperture centered on the pulsar, restricting the energy range of extracted photons to 3–10 keV for GIS and 3–12 keV for the other instruments. For the PCA data, in this energy band, we further restrict our search to Layer 1 data only, which provides the best sensitivity for a Crab-like spectrum; For the higher energy analysis afforded by the PCA, above ~ 12 keV, we used data from all three PCA layers.

We searched each data set for the expected 69 ms period predicted from the initial period and period derivative measurement (Torii *et al.* 1998). A periodogram was constructed using the χ^2 statistic to test against a null hypothesis. For each trial period, we folded the data into 10 bins and computed the χ^2 of the resultant profile. We search a narrow range of periods centered on the expected period ± 0.1 ms, sampled in increments of $0.1 \times P^2/T$, where T is the observation duration, and P is the test period. A highly significant signal was detected from each of our data sets.

As well as the newly obtained data, we have re-analyzed the previous GINGA and ASCA data (Aoki *et al.* 1992; Torii *et al.* 1998) in a uniform way and revised the period and its error by using the method of Leahy (1987). Our X-ray timing results derived from these 13 measurements are listed in table 1, along with an updated radio ephemeris (Kaspi 1999, Private communication).

3.2. Spectrum

We search for spectral dependence of the pulse profile by comparing the folded light curves in several energy bands. No strong energy dependence is evident in the energy resolved light curves. Furthermore, the pulse amplitude and pulse profile remained unchanged between observational epochs.

We examined the ASCA and SAX data on AXS J161730–505505 for any long term changes in its energy spectrum or flux. As for the timing analysis, we restrict our comparison to the energy range above 3 keV and extract photons from an 8' diameter aperture centered on the source. We fitted the spectrum with a power law function modified by interstellar absorption. The absorption was fixed at $6.8 \times 10^{22} \text{ cm}^{-2}$ (Torii *et al.* 1998). Spectra from each observation were found to be consistent with each other. Com-

binning the 7 ASCA observations, we obtain the pulse phase averaged photon index, $\Gamma = 1.4 \pm 0.2$, and the observed flux of $(3.6 \pm 0.2) \times 10^{-12} \text{ erg s}^{-1} \text{ cm}^{-2}$ (90% confidence errors) in the 3–10 keV range, which is consistent with the previous measurement (Torii *et al.* 1998).

4. DISCUSSION

A χ^2 fit to the all 14 data points as summarized in table 1 gives the mean spin-down rate of $\dot{P} = 1.3611(1) \times 10^{-13} \text{ s/s}$ and $P = 0.069347150(1)$ at MJD 50,000.0 (Figure 1). However, the quality of the fit is bad, with $\chi^2/d.o.f. = 4808/12$. The residual of the fit shows a jump of $\Delta P \simeq -1.2 \times 10^{-7} \text{ s}$ between the observations of 1993 August (MJD 49,217.6) and 1997 September (MJD 50,696.0) (Figure 2, top panel). Within the observation span of 10 years, the residual is neither periodic nor smooth. A sudden change in the period between MJD 49,217.6 and MJD 50,696.0 is suggested. We consider the most likely explanation for these residuals is due to a glitch, similar to those observed in several young rotation-powered pulsars.

Given the clear evidence of glitch activity, we next attempted to model the spin-down data with a single glitch followed by an exponential recovery. The data coverage is limited and we simply assumed the following relation for the spin-down.

$$P(t) = P_0 + \dot{P} \cdot (t - t_0) + \Delta P \cdot \exp\left(-\frac{t - t_0}{\tau}\right) \quad (1)$$

where $\Delta P = 0$ for $t < t_0$ and ΔP is a negative constant for $t \geq t_0$. This model contains five parameters, which, except for the depth of the glitch ΔP , are found to be independent of the time of the glitch, t_0 . The derived parameters, P_0 , $\dot{P}$, ΔP , and τ are summarized in table 2 for assumed values of $t_0 = 49,300.0$ MJD, $t_0 = 50,000.0$ MJD, and $t_0 = 50,600.0$ MJD. The residual for $t_0 = 50,000$ is shown in the middle panel of figure 2. The quality of the fit is now characterized by $\chi^2/d.o.f. = 22.6/10$.

The size of the glitch depends strongly upon the unknown glitch epoch t_0 . The fractional increase in rotation was found to be $\Delta P/P = -4.2 \times 10^{-6}$ for $t_0 = 50,000.0$ MJD (fixed) while it changes between $\Delta P/P = -11 \times 10^{-6}$ for $t_0 = 49,300.0$ MJD (fixed) and $\Delta P/P = -1.8 \times 10^{-6}$ for $t_0 = 50,600.0$ MJD (fixed). The minimum fractional increase in rotation rate is therefore comparable with those of the largest known pulsar glitches (Lyne, *et al.* 1996a; Shemar & Lyne 1996).

Using the above model, the recovery time following the glitch episode is found to be $\tau = 700$ days. This duration is somewhat unusual compared to the radio pulsars whose recovery time is seen to bifurcate between $\tau \simeq 100$ days and $\tau \geq 1,000$ days (Shemar & Lyne 1996). Our derived value lies squarely between these two time scales, perhaps due to the simple model we invoked which allows for only a single glitch. Because of the sparse data coverage between August 1993 and September 1997, however, we cannot determine if the recovery time could be expressed as the sum of the two timescales.

Glitches found in radio pulsars may be classified into three groups (Lyne & Graham-Smith 1998). The first is a Crab-like glitch which is characterized by steps mainly in $\dot{P}$. The second is a Vela-like glitch which is characterized by large changes in P ($\Delta P/P \simeq -10^{-6}$) and the exponential recoveries. For the Vela pulsar, linear sawtooth changes in $\dot{P}$ have been observed between glitches (Lyne *et al.* 1996b). The third kind is often found in old pulsars which is characterized by a change in P .

Since the large glitch found for AXS J161730–505505 is similar in its magnitude to those in the Vela pulsar, a phe-

nomenological model taking into account the sawtooth behavior may be a good description. Apart from short term effects, the spin-down for the Vela pulsar is expressed by a linear change in spin-down rate. Therefore, the following function is appropriate if the transient effects have already ceased by MJD 50,696.

$$P(t) = P_0 + \Delta P_0 + (\dot{P} + \Delta \dot{P}) \cdot (t - t_0) + \frac{1}{2} \ddot{P} \cdot (t - t_0)^2 \quad (2)$$

Here, $\Delta P_0 = 0$ and $\Delta \dot{P} = 0$ for $t < t_0$ and they are constant values ($\Delta P_0 < 0$ and $\Delta \dot{P} > 0$) for $t \geq t_0$. This model contains six parameters. Again, the time of the glitch, t_0 , had to be assumed. The derived parameters are summarized in table 3. For the condition that $\Delta P_0 < 0$, t_0 was restricted to $t_0 \lesssim 50,205$. The residual for $t_0 = 50,000$ is shown in the bottom panel of figure 2. The quality of the fit is significantly improved to $\chi^2/d.o.f. = 17.1/9$. Compared to the fit to Equation (1), the f-test gives a chance probability of ~ 0.1 . This result suggests that transient effects had indeed ceased by MJD 50,696. The reduced χ^2 is still larger than unity, suggesting the presence of timing noise and smaller glitches.

We can estimate the expectancy of large ($|\Delta P/P| > 10^{-7}$) glitches by using the semi-empirical relation based upon the superfluid vortex unpinning model (Alpar & Baykal 1994). Using this relation, the expected number of large glitches between the first *GINGA* observation and the last *ASCA* observations is 3.7. Therefore, there should have been about 4 glitches of $|\Delta P/P| > 10^{-7}$. Indeed the residuals to the fit of Equation (1) or (2) still hint a small jump of $\Delta P = -5 \times 10^{-9}$ s between MJD 51,263.8 (1999 Mar.) and MJD 51,394.2 (1999

Aug). This may be another glitch of $\Delta P/P = -7 \times 10^{-8}$, much smaller in magnitude than the one near MJD 50,000, but still relatively large compared to those seen for most radio pulsars (Shemar & Lyne 1996). We have thus found a giant glitch of $|\Delta P/P| \gtrsim 10^{-6}$ at $t_0 \simeq 50,000$ MJD ($49,218 \leq t_0 \leq 50,696$) and possibly a glitch of $|\Delta P/P| \simeq 7 \times 10^{-8}$ at $t_0 \simeq 51,300$ MJD ($51,264 \leq t_0 \leq 51,394$).

The detection of a giant glitch from AXSJ161730–505505 gives a rare sample for studying the origin of pulsar glitches and the interior structure of neutron stars. In this context, regular timing observation of this pulsar in the radio band is quite important to monitor the onset of a glitch and following transient effects. Long term timing observations for measuring the braking index are desired. Searches for the highly absorbed X-ray emission from the supernova remnant associated with this pulsar should be undertaken with the next generation X-ray observatories.

Acknowledgments —

We thank Jules Halpern for a critical reading of the manuscript, the referee Andrew Lyne for invaluable comments and suggestions on the original draft, and Victoria Kaspi for communicating the revised radio ephemeris. Special thanks to Alan Smale for expediting the delivery of the *XTE* data and to the XTEhelp@athena.gsfc.nasa.gov crew for their patience and assistance. Part of this research has made use of data obtained through the HEASARC data center, provided by NASA/GSFC. E. V. G. and G. V. acknowledge the supported of NASA LTSA grant NAG 5-7935.

REFERENCES

- Alpar, M.A., & Baykal, A. 1994, *MNRAS*, 269, 849
 Anderson, P.W., & Itoh, N. 1975, *Nature*, 256, 25
 Aoki, T., Dotani, T., & Mitsuda, K. 1992, *IAU Circ.*, 5588
 Boella, G., Butler, R.C., Perola, G.C., Piro, L., Scarsi, L., & Bleeker, J.A.M. 1997, *A&AS*, 122, 299
 Bradt, H.V., Rothschild, R.E., & Swank, J.H. 1993, *A&AS*, 97, 355
 Gotthelf, E.V., Petre, R., & Hwang, U. 1997, *ApJ*, 487, L175
 Gotthelf, E.V., Petre, R., & Vasisht, G. 1999, *ApJ*, 514, L107
 Kaspi, V.M., Crawford, F., Manchester, R.N., Lyne, A.G., Camilo, F., D'amico, N., & Gaensler, B.M. 1998, *ApJ*, 503, L161
 Leahy, D.A. 1987, *A&A*, 180, 275
 Lyne, A.G., Kaspi, V.M., Bailes, M., Manchester, R.N., Taylor, H., & Arzoumanian, A. 1996a, *MNRAS*, 281, L14
 Lyne, A.G., Pritchard, Graham-Smith, F., & Camilo, F. 1996b, *Nature*, 381, 497
 Lyne, A.G., & Graham-Smith, F. 1998, *Pulsar Astronomy Second Edition* (Cambridge: Cambridge Univ. Press)
 Rots, A.H., et al. 1998, *ApJ*, 501, 749
 Saito, Y., et al. 1997, *ASCA News*, 5
 Shemar, S.L., & Lyne, A.G. 1996, *MNRAS*, 282, 677
 Tanaka, Y., Inoue, H., & Holt, S.S. 1994, *PASJ*, 46, L37
 Torii, K., Kinugasa, K., Toneri, T., Asanuma, T., Tsunemi, H., Dotani, T., & Mitsuda, K. 1998, *ApJ*, 494, L207
 Urama, J.O., & Okeke, P.N. 1999, *MNRAS*, 310, 313

TABLE 1
TIMING DATA FOR AXS J161730–505505.

Number	Satellite/ Observatory	Instrument	Obs Date ^a	Exposure (s)	Epoch ^b (MJD)	Period ^c (ms)
1	GINGA	LAC	1989 Mar 4	14,400	47590.2	69.31890(3)
2	ASCA	GIS	1993 Aug 17	35,353	49217.277625	69.338019(2)
3	ASCA	GIS	1997 Sep 4	58,063	50696.736757	69.355301(1)
4	XTE	PCA	1998 Jan 2	33,712	50816.132391	69.356717(1)
5	Parkes	64 m	1998 Jan 15	—	50829.7	69.356889(7) ^d
6	SAX	MECS	1998 Sep 17	16,288	51073.892337	69.359778(1)
7	ASCA	GIS	1999 Feb 12	15,405	51222.279184	69.361532(4)
8	ASCA	GIS	1999 Feb 20	16,232	51229.708310	69.361624(7)
9	XTE	PCA	1999 Mar 5	37,253	51243.247065	69.3617824(5)
10	ASCA	GIS	1999 Mar 6	20,700	51243.354377	69.361782(8)
11	SAX	MECS	1999 Mar 23	38,197	51261.461543	69.362000(3)
12	ASCA	GIS	1999 Mar 26	18,914	51263.519288	69.362021(3)
13	SAX	MECS	1999 Aug 4	53,115	51394.834928	69.363568(2)
14	ASCA	GIS	1999 Sep 13	17,796	51434.716949	69.364037(6)

^aStart time of the observation.

^bMiddle time of the observation except for the radio ephemeris.

^c1 σ error of the last significant figure is shown in parentheses.

^dThe radio ephemeris reported here (Kaspi 1999, Private communication) has been revised from that given in Kaspi *et al.* (1998).

TABLE 2
BEST-FIT PARAMETERS OF A GIANT GLITCH IN AXS J161730–505505 FOR
EQUATION (1).

Parameter	Values		
t_0 [MJD]	49,300 (fixed)	50,000 (fixed)	50,600 (fixed)
P_0 [s] ^a	0.069338991(2)	0.069347220(3)	0.069354272(5)
$\dot{P}$ [s/s]	$1.3605(5) \times 10^{-13}$	$1.3605(5) \times 10^{-13}$	$1.3605(5) \times 10^{-13}$
ΔP [s]	$-8(2) \times 10^{-7}$	$-2.9(2) \times 10^{-7}$	$-1.26(4) \times 10^{-7}$
τ [days]	$7(1) \times 10^2$	$7(1) \times 10^2$	$7(1) \times 10^2$

^aThe period just before the glitch at t_0 .

TABLE 3
BEST-FIT PARAMETERS OF A GIANT GLITCH IN AXS J161730–505505 FOR
EQUATION (2).

Parameter	Values		
t_0 [MJD]	49,300 (fixed)	50,000 (fixed)	50,205 (fixed)
P_0 [s]	0.069338980(3)	0.06934707(3)	0.06934943(3)
ΔP_0 [s]	$-4.6(3) \times 10^{-7}$	$-1.0(2) \times 10^{-8}$	$-3.1(3) \times 10^{-10}$
$\dot{P}$ [s/s]	$1.344(3) \times 10^{-13}$	$1.332(5) \times 10^{-13}$	$1.328(5) \times 10^{-13}$
$\Delta \dot{P}$ [s/s]	$5.8(7) \times 10^{-15}$	$5.8(7) \times 10^{-15}$	$5.8(7) \times 10^{-15}$
$\ddot{P}$ [s $^{-1}$]	$-2.0(3) \times 10^{-23}$	$-2.0(3) \times 10^{-23}$	$-2.0(3) \times 10^{-23}$




FIG. 1.— The pulse period evolution of AXS J161730–505505 over 10 years.




FIG. 2.— *Top Panel*) Pulse period residual as fitted by a linear function. *Middle Panel*) Pulse period residual as fitted to Equation (1) with a glitch at MJD 50,000. *Bottom Panel*) Pulse period residual as fitted to Equation (2) with a glitch at MJD 50,000.